

RESEARCH ARTICLE

Dietary Vitamin C and Lipid Feed Supplementations Influence the Expression of Immune-Related Genes in *Litopenaeus vannamei* (Boone 1931) Subjected to Thermal Stress

Eljenmer L. Sindo^{1,2} , Angel Queene D. Dequito^{2,3*} , Rex Ferdinand M. Traifalgar² 

¹Institute of Fish Processing Technology, College of Fisheries and Ocean Sciences, University of the Philippines Visayas, Miagao 5023, Iloilo, Philippines

²Institute of Aquaculture, College of Fisheries and Ocean Sciences, University of the Philippines Visayas, Miagao 5023, Iloilo, Philippines

³Institute of Bioinformatics and Systems Biology, College of Engineering Bioscience, National Yang Ming Chiao Tung University, Hsinchu, Taiwan

ABSTRACT

This study investigated the effects of dietary vitamin C (vitC) and lipid on the expression of immune-related genes in *Litopenaeus vannamei* postlarvae subjected to thermal stress, with water temperature fluctuating from 27.0 ± 1.1 to 34.0 ± 1.8 °C for 15 days. Three experimental diets were formulated containing two different levels of vitC (1 and 3 g kg⁻¹) and lipid (17 and 34 g kg⁻¹). Significantly higher total hemocyte count ($9.63 \times 10^5 \pm 3.28 \times 10^5$ cells ml⁻¹) was observed in shrimp fed 3 g kg⁻¹ vitC and 34 g kg⁻¹ lipid diets. Using real-time quantitative polymerase chain reaction and the comparative 2^{-ΔΔC_t} method, the dietary vitC level of 3 g kg⁻¹ significantly upregulated heat shock protein 90 (HSP90), heat shock protein 70 (HSP70), anti-lipopopolysaccharide factor (ALF), crustin, and the Prophenoloxidase (proPO) system. Meanwhile, the dietary lipid level of 34 g kg⁻¹ resulted in a significant upregulation of HSP90 and penaeidin 2 (PEN 2) genes. Regarding the growth and survival indices of the shrimps fed the formulated diets, no significant differences ($P > 0.05$) were observed. In conclusion, increased dietary supplementation of vitC (3 g kg⁻¹ diet) and lipid (34 g kg⁻¹ diet) upregulated the immune-related genes, enhancing the immunity of shrimps during thermal stress.

*Corresponding Author: addequito@up.edu.ph

Received: May 28, 2023

Accepted: December 11, 2024

Keywords: Pacific white shrimp, dietary supplementation, vitamin C, lipid, fluctuating temperature, immunity

1. INTRODUCTION

The Pacific white shrimp, *Litopenaeus vannamei* (Boone 1931), is among the widely cultivated fishery products and is a vital economic, food, and protein source (Fachri et al. 2021). From 1995 to 2019, it was the most significant contributor to global shrimp aquaculture production (FAO 2020), with an expected increase in inland culture due to its adaptability to low-salinity environments (Liao and Chien 2011).

Temperature is a crucial water parameter in shrimp culture, directly affecting shrimp's physiological performance, growth, survival, metabolic rate, and immune response (Zhang et al.

2019). Elevated water temperature stresses shrimp, resulting in poor immunity and resistance to infection, decreased phagocytosis activity, phenoloxidase activity, and respiration activity (Cheng et al. 2005). Fluctuations between optimal and high-water temperatures increase the infectivity of the white spot syndrome virus (WSSV) in juvenile *L. vannamei* (Rahman et al. 2007), while cyclic temperature fluctuations decrease growth rate, food utilization efficiency, and hemolymph glucose content (Guo et al. 2010). In other aquatic organisms, such as the Mozambique tilapia (*Oreochromis mossambicus*) exposed to temperatures as high as 35 °C, a significant reduction in the white blood cell count was observed, weakening the immune defenses of the organism

(Ndong et al. 2007). Ectothermic animals exposed to thermal stress suffer from oxidative stress and a shift in redox state (Nakano et al. 2014). Thermal stress results in adaptive responses in some aquatic animals, including crustaceans. However, these responses can be lost or dysfunctional when the animal is exposed to extreme and prolonged stress, resulting in growth inhibition, reproductive failure, and poor disease resistance (Wendelaar-Bonga 1997).

Vitamin C (vitC) is essential for penaeid shrimps and aquatic animals. It improves stress resistance, growth performance, physiological function, and immunological response (Dawood and Koshio 2018). It helps in collagen synthesis by hydroxylating proline and lysine (Niu et al. 2009). Aside from its immunomodulatory properties, it is reported to have potent antioxidant properties that protect low-density lipoproteins (LDLs) from oxidation (Cheng et al. 2005). In *L. vannamei*, dietary vitC intake improves growth, survival, and antioxidant status (Ebadi et al. 2021). In genetically modified *O. niloticus*, increasing dietary vitC supplementation significantly increased weight gain, liver vitC content, muscle collagen, and total body lipid content (Huang et al. 2016).

Eicosapentaenoic (20:5n-3, EPA), docosahexaenoic (22:6n-3, DHA), and arachidonic (20:4n-6, ARA) acids are polyunsaturated fatty acids (PUFAs) essential for many physiological functions in shrimp, including survival, reproduction, growth, and immunological responses (Mercier et al. 2009). The n-3 PUFAs in dietary lipids, such as in fish oil, reportedly mediate the fluidity and restructuring of cellular membranes and alter the subcellular localization of proteins (Matesanz et al. 2010). EPA-rich fish oil is an alternative cyclooxygenase (COX) substrate and enhances lipid and lipoprotein metabolism. By increasing reduced glutathione (GSH) and HSP70, fish oil is also essential for reducing oxidative stress and minimizing cellular damage caused by heat stress (Millward 2012).

Recent studies have focused on examining immune-related genes and their expression levels as a secondary response to stress, including exposure to extreme temperatures. The family of Heat Shock Proteins (HSPs), such as HSP90 and HSP70, are known for their immediate response to stressors (Junprung et al. 2021). These immune genes aid in refolding, folding, translocating, assembling, and destroying proteins (Junprung et al. 2021) when animals are in stressful conditions (i.e., temperature stress, toxic stress, and infection). The Prophenoloxidase (proPO) system functions in tissue repair and defense against pathogens (Junprung et al. 2021). Penaeid shrimp

also rely on antimicrobial peptides (AMPs) against invading pathogens (Destoumieux-Garzón et al. 2016). The AMPs found on shrimp hemocytes, such as penaeidin 3 (PEN 3) and penaeidin 2 (PEN 2), reportedly have antibacterial and antifungal activities in response to microbial infection (Destoumieux-Garzón et al. 2016). Crustin is a polypeptide with antiprotease properties, while anti-lipopolysaccharide factors (ALFs) have antifungal, antibacterial, and immunomodulatory abilities (de la Vega et al. 2008). Thermal stress weakens the pathogen resistance and immune response of shrimps (Rahman et al. 2007; Janewanthanakul et al. 2020). During these conditions, AMPs may provide immune defense due to their immunomodulatory and antibacterial properties. Pathogens can also cause tissue damage, and proPO may mediate these effects due to their role in repairing damaged tissue.

This study aimed to investigate the effects of increased dietary supplementation levels of vitC and lipid on *L. vannamei* postlarvae subjected to thermal stress, specifically on the expression levels of immune-related genes: HSP70, HSP90, ALF, PEN 3, PEN 2, Crustin, and proPO, with EF1a as the reference gene. The study also investigated its impacts on the total hemocyte count (THC), growth indices (specific growth rate, weight gain, and feed conversion ratio), and survival rate.

2. MATERIALS AND METHODS

2.1 Experimental animal and acclimation

Postlarvae (PL 12) specific pathogen-free (SPF) Pacific white shrimps, *L. vannamei* (Boone 1931), were purchased from HP Aquafarm Inc. Hatchery (Iloilo, Philippines) and transported to the Institute of Aquaculture Multi-Species Hatchery Complex, University of the Philippines Visayas. Prior to the experiment, randomly selected SPF shrimps were acclimated for seven (7) days in a 10-ton concrete tank. The seawater (35‰) was treated with 10 ppm chlorine. After 24 h, the seawater was neutralized with the same amount of thiosulfate prior to stocking, following the manufacturer's instructions (Rainbow™ OTO Solution Chlorine/Bromine Test). The rearing conditions were maintained at a temperature set at 27.5 ± 0.5 °C with moderate aeration and natural photoperiod. The acclimated shrimps were hand-fed with commercial feed (Vannamei #3 starter pellets, Oversea Aqua Feeds, Philippines, 37% crude protein) at 5% of their biomass daily at 0800, 1200, and 1600 h.

2.2 Experimental diet

Three experimental diets in sinking pellet form were formulated to contain different amounts of vitC (ascorbic acid monophosphate) and dietary lipid (Danish fish oil). The Danish fish oil has around 1–7 % free fatty acids sourced from fresh fish (FE, Skagen). The diets were prepared following Genio et al. (2015) and Traifalgar et al. (2010). VitC and lipid are crucial inclusions in the practical and basal diets of shrimps for growth, development, and immune responses (Montayo and Molina 1995; Gonzalez-Felix et al. 2002; Lee and Shiau 2002). The dietary supplementation level in Treatment 1 (control) for vitC was from Montayo and Molina (1995) and Ebadi et al. (2021). In the same treatment, the dietary supplementation level of lipids was within the practical range provided by Gong et al. (2001), Niu et al. (2011), and Mawalat et al. (2016). For 15 days, shrimp were fed the formulated experimental diets at 5% of their biomass, subdivided

into three feeding rations at 0800, 1200, and 1600 h. Treatment 1 (vitC+lipid) contained 1 g kg⁻¹ vitC and 17 g kg⁻¹ fish oil and was used as the control. Treatment 2 (3vitC+lipid) contained 3 g kg⁻¹ vitC and 17 g kg⁻¹ fish oil. Treatment 3 (vitC+2lipid) contained 1 g kg⁻¹ vitC and 34 g kg⁻¹ fish oil. The ingredients and the proximate composition of the formulated experimental feeds are presented in Table 1.

2.3 Experimental design

One hundred eighty (180) *L. vannamei* (4.0 ± 1.0 g) were randomly distributed among nine 120-L rectangular plastic tanks. Each treatment was in triplicate, resulting in nine tanks. Seawater was subjected to the same treatment method done during the acclimation process. Shrimps were subjected to thermal stress using an electric heater with a built-in thermometer turned on and off daily at 0700 and 1600 h, respectively, simulating fluctuating environmental

Table 1. Ingredients, formulation, and proximate composition of the experimental diets vitC+lipid, 3vitC+lipid, and vitC+2lipid.

Ingredients	Diets		
	vitC+lipid	3vitC+lipid	vitC+2lipid
Soybean meal (48 solv)	230	230	230
Trace mineral premix ^a	15	14	15
Squid meal	80	80	80
Fish meal (70 % - Danish)	230	230	230
Rice bran	125	125	125
Corn/wheat starch	240	240	240
Fish oil	17	17	34
Soybean oil	17	17	0
Vitamin premix ^b	15	14	15
Lecithin - Soy (70 %)	10	10	10
Gluten (wheat)	20	20	20
Vitamin C ^c	1	3	1
Total:	1000	1000	1000
Proximate composition			
Crude Protein	37.5 ± 0.17	35.8 ± 0.08	37.1 ± 0.02
Crude lipid	6.4 ± 0.16	10.6 ± 0.12	7.6 ± 0.04
Ash	7.95 ± 0.04	7.63 ± 0.06	8.03 ± 0.06
Moisture	7.4 ± 0.01	7.2 ± 0.02	7.0 ± 0.02
Crude Fiber	2.9 ± 0.02	2.9 ± 0.05	2.9 ± 0.09
Nitrogen-Free Extract	37.8 ± 0.06	35.8 ± 0.03	37.5 ± 0.03

^aTrace mineral premix (g kg⁻¹ in diet): CaHPO₄ · 2H₂O, 8 g; MgSO₄ · 7H₂O, 5 g; KH₂PO₄, 4 g; Na₂H₂PO₄ · 2H₂O, 2 g; MnSO₄ · H₂O, 0.6 g; FeSO₄, 0.6 g; ZnSO₄ · 7H₂O, 0.6 g; CO(NO₃)₃, 0.1 g; CuSO₄ · 5H₂O, 0.1 g.

^bVitamin premix (mg, mcg or IU kg⁻¹ in diet): vit. A, 1200000 IU; vit. D₃, 200000 IU; vit. E, 20000 IU; vit. B₁, 8000 mg; vit. B₂, 8000 mg; vit. B₆, 5000 mg, vit. B₁₂ (1 %), 2000 mcg; Niacin, 40000 mg; Calcium Pathothenate, 20000 mg; Biotin, 40 mg, Folic Acid, 1800 mg; Ethoxyquin, 500 mg.

^c35% min ascorbic acid monophosphatet

temperatures. Shrimps were fed the experimental diets 1 h after being subjected to thermal stress. The recorded temperature in all tanks fluctuated from 27.0 ± 1.1 to 34.0 ± 1.8 °C. Shrimp exposed to temperatures above 34 °C are considered stressed (Allan et al. 2006), which adversely affects optimal shrimp growth performance and increases the occurrence of diseases in shrimp culture (Alonzo et al. 2017). In each tank, 80% of the seawater volume was siphoned off and replaced with treated seawater weekly. The water temperature and salinity were measured at 0800 and 1400 h. Temperature was determined using an alcohol capillary laboratory thermometer, while salinity was measured using the REB-32ATC 0–32 % Brix refractometer.

2.4 Sampling

Random sampling was done on days 0, 7, and 15 for growth indices, THC, and gene expression analysis. Ribonucleic acid (RNA) samples were extracted from the hemolymph of three shrimp per tank (9 shrimp per treatment) and pooled in one (1) tube as a sample. Average body weight (ABW) was determined randomly from five to ten individuals per tank. Weight gain rate (WGR), specific growth rate (SGR), feed conversion ratio (FCR), and survival rate (SR) were calculated following Genio et al. (2015).

2.5 Histological examination

Using Davidson's alcohol-formalin-acetic acid (AFA), hepatopancreas and gill tissues of *L. vannamei* were fixed for 48 hours and transferred to 70 % ethanol (Bell and Lightner 1988). After dehydration in increasing ethanol concentrations, the tissue samples were cleared with xylene and embedded in paraffin. Microtome-cut sections (5µm-thick) were stained using Hematoxylin and eosin (H&E). The histological slide sections were examined using light microscopy.

2.6 Total hemocyte count (THC) determination

Three shrimp individuals per treatment were collected for hemocyte count analysis during each sampling period. Precooled 0.4 mL anticoagulant composed of shrimp salt solution (SSS) containing 450 mM NaCl, 10 mM HEPES, 10 mM EDTA, and 10 mM KCl at pH 7.3 and 850 mOsm kg⁻¹ (Hernandez-Lopez et al. 1996) was prepared in each syringe. A 1 mL sterile syringe with a 25-gauge needle was used to extract 0.2 mL of hemolymph from the first abdominal

segment close to the genital pore of the shrimp. The hemolymph samples were then transferred to a 1.5 mL microcentrifuge tube containing a fixative solution (10% formalin in 0.45 M NaCl) and kept at 4°C. The fixed sample was added with a drop of Rose Bengal solution (1.2% Rose Bengal in 50% ethyl alcohol) and then stained for 20 minutes, following Genio et al. (2015). Hemocytes were counted using the Neubauer hemocytometer, and the total hemocyte count values were expressed as cells ml⁻¹, calculated following Joseph and Philip (2007).

2.7 RNA extraction

Hemolymph samples were extracted from the pleopod base at the first abdominal segment using a 1 mL syringe fitted with a 25-gauge needle. The extracted hemolymph was diluted with an anticoagulant medium prepared according to Hernández-López et al. (1996). The anticoagulant-hemolymph mixture was centrifuged at 4 °C at 10000 × g for 5 minutes, with the plasma drained after centrifugation. The cell pellet was added with 500 Trizol (Invitrogen) and stored at -20 °C.

2.8 Immune-gene expression analysis by qRT-PCR

The concentration and quality of the extracted RNA were assessed using a spectrophotometer, 1% agarose gel electrophoresis, and Qubit (Invitrogen). The SuperScript™ First-Strand Synthesis System test kit from Thermo Fisher Scientific was used to synthesize cDNA following the manufacturer's procedure. The qRT-PCR gene analysis was performed in CFX Connect Real-Time PCR (Bio-Rad) and iQ SYBR Green Supermix (Bio-Rad). The forward and reverse sequences of the nucleotide primers are shown in Table 2. The qPCR analyzed data were interpreted using the comparative 2^{-ΔΔCt} method (Livak and Schmittgen 2001) to compute the relative mean fold expression of the target genes. Fold differences and cycle threshold (Ct) values were recorded using Cycling Fluorescence Excitation Manager software (Bio-Rad). Gene expression values are presented as geometric means.

2.9 Proximate analysis

The formulated feed was evaluated using the following methods and analytical tools: crude protein using the Kjeldahl method with block digestion and steam distillation (FOSS Tecator Digestion and

Table 2. Nucleotide sequence of the primers (HSP70, HSP90, Crustin, PEN 3, Crustin, and ALF, PEN 2, proPO) with EF1 as the reference gene.

Primer	Sequence (5' – 3')	References
LvHSP70-F	CTCCTGCGTGGGTGTGTT	Janewanthanakul et al. (2020)
LvHSP70-R	GCGGCGTCACCAATCAGA	Janewanthanakul et al. (2020)
LvHSP90-F	TGGGCTTCTACTCCGCCTACC	Junprung et al. (2017)
LvHSP90-R	ACGGTGAAAGAGCCTCCAGCA	Junprung et al. (2017)
LvPEN3-F	CACCCTTCGTGAGACCTTTG	Junprung et al. (2019)
LvPEN3-R	AATATCCCTTTCCCACGTGAC	Junprung et al. (2019)
LvALF-F	GTTATCACGCCCTTTTCTAC	Junprung et al. (2017)
LvALF-R	GAAGAATGACCTGTCCAAC	Junprung et al. (2017)
LvCrustinI-F	GCTGGCCTCGATAAGTGTTG	Janewanthanakul et al. (2020)
LvCrustinI-R	CATCGGTCTTCTTCAGATG	Janewanthanakul et al. (2020)
LvPEN2-F	TCGTGGTCTGCCTGGTCTT	Junprung et al. (2019)
LvPEN2-R	CAGGTCTGAACGGTGGTCTTC	Junprung et al. (2019)
LvproPO1-F	AACTCCATTCCGTCCGTCTG	Junprung et al. (2019)
LvproPO1-R	GGCTTCGCTCTGGTTAGGAT	Junprung et al. (2019)
LvEF1a-F	CTTGATTGCCACACTGCTCAC	Janewanthanakul et al. (2020)
LvEF1a-R	TCTCCACGCACATAGGCTTG	Janewanthanakul et al. (2020)

FOSS Kjeltex 8200 Auto Distillation Unit), crude fat was determined using the Soxtec extraction method (FOSS Soxtec 2050 Automatic System), crude fiber determined using the AOAC Ceramic Fiber Filter method (FOSS Fibertec 2010 System), moisture content was determined using the Mettler Tornado Halogen Moisture Analyzer, ash was determined using the AOAC method (2010) and the acid value was generated using the by Soxtec extraction or titrimetric method.

2.10 Data and statistical analysis

All analyzed data are presented as mean \pm standard errors (SE; $n = 3$). The data were within normal distribution as verified by the Shapiro-Wilk test. The one-way analysis of variance (ANOVA) was used to examine significant differences, $P < 0.05$.

When significant differences were detected, the nature of these differences was determined using Tukey's test.

3. RESULTS

3.1 Growth indices and survival rate

The growth and survival indices of the shrimp subjected to thermal stress and fed three distinct formulated diets are presented in Table 3. The formulated diets contained 1 g kg⁻¹ vitC and 17 g kg⁻¹ lipid (vitC+lipid), 3 g kg⁻¹ vitC and 17 g kg⁻¹ lipid (3vitC+lipid), and 1 g kg⁻¹ vitC and 34 g kg⁻¹ lipid (vitC+2lipid).

When subjected to thermal stress, shrimp fed vitC+2lipid resulted in the highest survival rate of 81.67 \pm 7.26 and the lowest FCR of 1.41 \pm 0.11. Meanwhile, 3vitC+lipid resulted in the highest SGR

Table 3. Growth indices and survival rate of *L. vannamei* ($n = 20$) subjected to thermal stress.

Parameters	vitC+lipid	3vitC+lipid	vitC+2lipid
Initial weight (g)	4.07 \pm 0.05	4.08 \pm 0.05	4.10 \pm 0.19
Final weight (g)	8.52 \pm 0.24	8.96 \pm 0.50	8.55 \pm 1.12
Weight gain rate (%)	209.49 \pm 3.39 ^a	219.32 \pm 9.30 ^a	208.47 \pm 24.89 ^a
Specific growth rate (%)	4.60 \pm 0.10 ^a	4.90 \pm 0.26 ^a	4.50 \pm 0.79 ^a
Survival rate (%)	71.67 \pm 4.41 ^a	76.67 \pm 4.41 ^a	81.67 \pm 7.26 ^a
Feed conversion ratio (FCR)	2.08 \pm 0.44 ^a	1.60 \pm 0.33 ^a	1.41 \pm 0.11 ^a

Shrimp fed diets with 1 g kg⁻¹ vitC and 17 g kg⁻¹ lipid (vitC+lipid), 3 g kg⁻¹ vitC and 17 g kg⁻¹ lipid (3vitC+lipid), and 1 g kg⁻¹ vitC and 34 g kg⁻¹ lipid (vitC+2lipid). Data are mean \pm SE. Different superscripts indicate significant differences ($P < 0.05$).

of 4.90 ± 0.26 and WGR of 219.32 ± 9.30 . Shrimp fed vitC+lipid resulted in the lowest survival rate of 71.67 ± 4.41 and the highest FCR of 2.08 ± 0.4 . No significant differences were observed between diets ($P > 0.05$).




Figure 1. THC of *L. vannamei* ($n = 20$) subjected to thermal stress fed diets with 1 g kg^{-1} vitC and 17 g kg^{-1} lipid (vitC+lipid), 3 g kg^{-1} vitC and 17 g kg^{-1} lipid (3vitC+lipid), and 1 g kg^{-1} vitC and 34 g kg^{-1} lipid (vitC+2lipid). Data are mean \pm SE. Different superscript on Day 7 and 15 sampling period indicates significant differences among groups ($P < 0.05$).




Figure 2. Mean fold gene expressions of heat shock protein70 (HSP70) gene, heat shock protein90 (HSP90) gene, anti-lipopolysaccharide factor (ALF) gene, penaeidin 3 (PEN 3) gene, penaeidin 2 (PEN 2) gene, crustin, and prophenoloxidase (proPO) system gene measured at (A) day 7 and (B) day 15 of culture from the hemolymph of *L. vannamei* subjected to thermal stress, fed vitC and lipid-supplemented diets (vitC+lipid, 3vitC+lipid, and vitC+2lipid). Gene expressions are presented in geometric mean. Data are mean \pm SE. Different superscripts indicate significant differences ($P < 0.05$).

3.2 Nonspecific immune response, THC

The THC of *L. vannamei* subjected to thermal stress and fed vitC+lipid, 3vitC+lipid, and vitC+2lipid is shown in Figure 1. At day 7, vitC+2lipid resulted in a significantly higher ($P < 0.05$) THC of $7.88 \pm 3.23 \times 10^5$ cells ml^{-1} than vitC+lipid, which resulted in a THC of $4.71 \pm 2.89 \times 10^5$ cells ml^{-1} . Meanwhile, 3vitC+lipid resulted in THC of $7.07 \pm 5.55 \times 10^5$ cells ml^{-1} which is not significantly different ($P > 0.05$) from that of vitC+lipid and vitC+2lipid. At day 15, 3vitC+lipid resulted in a significantly lower ($P < 0.05$) THC of $5.44 \pm 1.84 \times 10^5$ cells ml^{-1} than vitC+lipid and vitC+2lipid with THC of $9.63 \pm 3.28 \times 10^5$ cells ml^{-1} and $9.98 \pm 5.29 \times 10^5$ cells ml^{-1} respectively.

3.3 Expression of immune-related genes in *L. vannamei*

The mean fold expressions of the immune-related genes at days 7 and 15 from the hemolymph samples of *L. vannamei* subjected to thermal stress are illustrated in Figure 2. Compared with the vitC+lipid diet, the vitC+2lipid diet at day 7 of culture resulted in a significantly upregulated HSP90 and PEN 2 ($P < 0.05$) at 2.16-fold and 1.69-fold, respectively. Notably, the vitC+2lipid diet resulted in a significantly downregulated Crustin ($P < 0.05$); however, it is not downregulated in shrimp fed with vitC+lipid or 3vitC+lipid diets. The experimental diets resulted in HSP70, ALF, PEN 3, and proPO levels that were not significantly expressed.

Immune-related genes in shrimp fed with the 3vitC+lipid diet that were not found to be significantly upregulated at day 7, such as ALF and HSP70, were observed to be significantly upregulated ($P < 0.05$) at day 15 of culture. Compared with the vitC+lipid diet, the 3vitC+lipid diet significantly resulted in a 4.33-fold higher ALF expression, 2.61-fold higher HSP70 expression, 1.83-fold higher HSP90 expression, and 1.64-fold higher proPO expression ($P <$

0.05). However, the vitC+2lipid diet downregulated the expressions of HSP70, HSP90, ALF, PEN3, PEN2, Crustin, and proPO.

4. DISCUSSION

Long-term hyperthermal stress above the critical thermal point disrupts normal physiological processes in shrimp. Extreme temperature conditions decrease food intake, impede molting and growth, and weaken the immune system of shrimps, leading to mortality (Roberts et al. 2010). Thermal stress also leads to increased incidence and prevalence of diseases in shrimp culture, such as susceptibility to WSSV and *Vibrio alginolyticus* (Cheng et al. 2005)

Vitamin C is a known immunostimulant, while lipids from fish oil reportedly have beneficial effects on the immunity of some aquatic organisms (Kumar et al. 2022). However, studies examining the effects of increased vitamin C and lipid intake on *L. vannamei* exposed to thermal stress are scarce. Therefore, the impact of these supplements on shrimp needs to be investigated to possibly improve the animals' immune response to thermal stress.

4.1 On growth indices and survival rate

The effects of increased dietary vitC and lipid supplementation on the growth and survival indices of *L. vannamei* exposed to thermal stress were not significantly different between groups ($P > 0.05$). However, shrimp fed diets with increased vitC and lipid supplementation had numerically better growth (i.e., weight gain, specific growth rate), survival, and feed conversion ratio. For comparison, *L. vannamei* reared at 25 ± 4 °C and fed only the optimal amounts of required nutrients already had significantly lower growth rates than shrimp reared at 25 ± 3 °C, 25 ± 1 °C, and 25 ± 1 °C (Guo et al. 2010). The physiological parameters of the shrimp at 25 ± 4 °C indicated a stressed state. Long-term hyperthermal stress affects the physiological processes of shrimp, resulting in stunted growth (Roberts et al. 2010). Water temperatures above 34 °C negatively affect the metabolic rate of *L. vannamei*, resulting in growth discrepancies and even retardation (Allan et al. 2006). In the present study, the growth and survival indices of *L. vannamei* postlarvae exposed to thermal stress and fed diets with increased vitC and lipid supplementation still showed better growth performance. These results may indicate that the increased supplementations promoted shrimp growth and survival. This is consistent with the results of Chien et al. (1999) with spiny dogfish *Terapon*

jarbua, as the test animal, reared at 36 °C. The best growth performance was observed in dogfish fed diets with increased vitC supplementation. Mortality also decreased with increasing vitC supplementation levels, even with increasing water temperature.

The proximate analysis showed a reduced crude protein content of the 3vitC+lipid diet. Despite this, shrimps fed 3vitC+lipid resulted in the highest WGR and SGR. The effect of vitamin C on crude protein content necessitates further studies, as it may be affected by different factors. For instance, the proximate composition of the muscle of *L. vannamei* fed different levels of dietary vitamin E, lipid, and vitamin C showed varying crude protein levels (Ebadi et al. 2021). When vitamin E supplementation is increased in diets with the same vitamin C levels, crude protein content decreases. Meanwhile, crude protein content increases when vitamin C levels are higher than vitamin E in the same diet formulation. The same study also noted that vitamin C and E interaction significantly affected the antioxidant activity and shrimp growth, thereby improving growth indices and reducing dietary lipid levels. Van Gaal and De Leeuw (1989) suggested that nutrient combustion rate changes, such as increased lipid combustion, can impact protein levels. Mrigal carp (*Cirrhinus mrigala*) fingerlings fed diets of varying vitamin C levels improved body protein content and reduced body fat, which indicates the more pronounced effect of vitamin C (Zehra and Khan 2012). Fracalossi et al. (1998) likewise reported the same differences in weight gain and protein content as a function of the vitamin C-free and enriched diets. In the present study, the increased vitamin C supplementation may not directly compromise protein content, but it can interact with other dietary components to influence protein content and metabolism.

4.2 On nonspecific immune response, THC

When the effect of vitC and lipids on the THC of *L. vannamei* postlarvae exposed to thermal stress was determined, shrimp supplemented with vitC and lipids had significantly higher THC. In contrast to these findings, Díaz et al. (2013) reported a significant THC decrease when *L. vannamei* were exposed to their critical thermal maximum (36–41.6 °C), while de la Vega et al. (2008) found no significant differences in THC when *Penaeus monodon* were exposed to 35 °C after 24 h. Thermal stress also significantly reduced the THC of *L. vannamei* after 6–12 h of rearing in freshwater or seawater (Jia et al. 2014). However, in these reports, the test shrimps were

only fed the required basal diet. The increased vitamin C and lipid supplementation in the present study may have contributed to the increased THC, even when the shrimps were subjected to thermal stress. In other aquatic species, increased supplementation of vitamin C at 250.12 to 321.38 mg kg⁻¹ in red swamp crayfish (*Procambarus clarkii*) diet resulted in increased THC, improved growth, immunity, and antioxidant capacity (Kong et al. 2021).

Vitamin C plays a key role in hemocyte (blood cell) formation due to its direct role in collagen synthesis (Niu et al. 2009). Collagen, a structural protein, is critical for the stability of blood vessels and, thus, for the proper functioning of the bone marrow, where red blood cells are formed (Mercier et al. 2012). The antioxidant properties of vitamin C also protect red blood cells by scavenging free radicals and minimizing oxidative stress (Cheng et al. 2005). Although the role of fish oil in hemocyte formation is not well established, the mediating effect of fish oil on cell membrane integrity (Matesanz et al. 2010) is hypothesized to have beneficial effects on hemocytes. The structural stability and functionality of cell membranes are influenced by fatty acids esterified in phospholipids. Robust cell membranes contribute to efficient oxygen transport in the blood, and omega-3 fatty acids have been shown to increase the ability of hemoglobin to transfer oxygen by making red blood cells more flexible and deformable (Patel et al. 2013). Increasing hemocyte count is critical, especially when these cells are also affected by environmental stressors such as pH, salinity, and temperature (Muthaiyan 2020). They are among the most critical components of crustacean cellular immune response, as they help remove foreign material by influencing cell adhesion and cytotoxicity and induce hemocyte-mediated defense responses (Cheng et al. 2005).

Interestingly, the THC of shrimp fed the 3vitC+lipid diet significantly decreased compared to those fed the vitC+lipid and the vitC+2lipid diets on day 15. This contrasts with the immediate reduction in THC observed by Cheng et al. (2005) and de la Vega et al. (2008). It could be that the beneficial effects of vitamin C peaked on day 7, and the shrimp had already acclimated to the increased vitamin C intake. THC could also be affected by treatment and timing of hemolymph collection. *L. vannamei*, infected with the infectious myonecrosis virus (IMNV), did not show an increase in THC until 17 days after the viral challenge ended (Sabry-Neto and Nunes 2015). This could help explain the (reduced) variability of THC in shrimp fed 3vitC+lipid compared with vitC+lipid only and vitC+2lipid, as it might be influenced by treatment

and sampling conditions. Pan et al. (2008) also noted that THC and the number of large granular cells could stabilize after six days without significant differences. In any case, studies on the long-term effects of vitamin C on the THC of shrimp exposed to thermal stress are recommended for a more conclusive impact of dietary supplementation.

4.3 On the expression of immune-related genes in *L. vannamei*

The effects of increased dietary vitC and lipid supplementation on the expression of the immune-related genes HSP70, HSP90, ALF, PEN 3, PEN 2, Crustin, and proPO of *L. vannamei* exposed to thermal stress were also investigated. Significant upregulation of target genes was found in shrimp with increased vitC and fish oil supplementation. Shrimp supplemented with vitC exhibited significant upregulations of HSP70, HSP90, ALF, Crustin, and proPO. Vitamin C is considered an immunostimulant due to its ability to enhance the nonspecific immune response of most crustaceans, which may explain the upregulations of the target genes observed in the present study. It is a cofactor and mediates enzymatic reactions and cytokine synthesis, regulating the immune responses in shrimp (Harrison and May 2009). In loaches, *Misgurnus anguillicaudatus* reared at 34 °C, a diet supplemented with 1000 mg kg⁻¹ vitC, which is five (5) times higher than their optimal vitamin C requirement showed significantly higher HSP70, HSP90 α , HSP90 β expressions (Yan et al. 2017). It is hypothesized that the increased vitamin C supplementation may have reduced the damage caused by thermal stress. Wuchang bream *Megalobrama amblycephala* under thermal stress (34 °C) fed with increased vitamin C supplementation resulted in better antioxidant activity, nonspecific immunity, and HSP70s mRNA expression, increasing thermal stress resistance (Ming et al. 2012). Ebadi et al. (2021) reported that increased vitamin C supplementation in the diet significantly improved survival, growth performance, and antioxidant activity of *L. vannamei* when the temperature was controlled and in the optimal range. The same conclusion was observed by Lee and Shiau (2002), where high ascorbate supplementation, regardless of source, significantly improved the immunological indices of *P. monodon* shrimp. These studies show that the beneficial effects of vitamin C are still observed when experimental animals are under thermal stress, as in the present study.

Fish oil is known for its beneficial effects

on nutrition, growth, and development because it is the primary source of dietary lipids rich in highly unsaturated fatty acids or HUFAs (Zhou et al. 2007). In the present study, shrimp fed vitC+2lipid showed significantly upregulated HSP90 and PEN2. Other studies have also reported that fish oil positively affects immune-related gene expressions in animals under thermal stress. Chim et al. (2001) showed that increased levels of HUFAs in the diet resulted in better adaptation and resistance of *Penaeus stylirostris* to environmental stress (heat and salt stress) by altering cellular membrane composition. A diet enriched with HUFAs also increases the lipid fraction of circulating lipoproteins (Vargas-Albores 1995). In agreement with these studies, it only shows that a fish oil-enriched diet of *L. vannamei* subjected to thermal stress positively affects its immune-related genes and, therefore, immune response.

5. CONCLUSION

Increased supplementation of dietary vitamin C and lipid on *L. vannamei* postlarvae subjected to thermal stress significantly upregulated the expression of immune-related genes HSP70, HSP90, proPO, and AMPs, compared to shrimp fed only the minimum basal diet and nutrient levels. The increased supplementation also resulted in numerically better growth and survival indices and feed conversion ratio. The findings suggest that farmers involved in *L. vannamei* culture may increase dietary vitC and lipid supplementation up to 3 g kg⁻¹ and 34 g kg⁻¹, respectively, to boost the immunity of the shrimp when exposed to thermal stress. The growth and survival of the shrimp are also unaffected by the increased supplementation levels. Nonetheless, an additional round of feeding experiments is recommended for short-term feeding trials to check and verify the growth and survival indices. Studies determining the long-term impact of further elevated levels of dietary vitamin C and lipid supplementations on shrimps subjected to thermal stress are recommended. The optimal profit generated when increased dietary vitamin C and lipid supplementations are added to the feed costs also calls for further studies.

ACKNOWLEDGMENT

The Department of Science and Technology - Philippine Council for Agriculture, Aquatic and Natural Resources Research and Development (DOST-PCAARRD) and the UP Visayas CFOS-IA

TGMPEFET Project are greatly appreciated by the authors for funding the research. The Department of Agriculture – Biotech Scholarship Program is also acknowledged for the undergraduate thesis grant. The authors recognize Ms. Rowena Cadiz, Ms. Mary Antonette Atilano, Ms. Emelyn Joy Mameloco, Ms. Ma. Delza Anduque, and Mr. Jeymarch Felongco for providing technical support.

AUTHOR CONTRIBUTIONS

Dequito A: Conceptualization, Experimentation, Analysis, Writing – Final Review and Editing, Funding Acquisition, Project Administration and Supervision. **Sindo E:** Conceptualization, Experimentation, Analysis, Writing – Original Draft. **Traifalgar R:** Conceptualization, Writing – Review and Editing, Project Supervision. Dequito A. and Sindo E. contributed equally to this work.

CONFLICTS OF INTEREST

The authors affirmed that the work provided in this report does not present any conflicts of interest.

ETHICS STATEMENT

The study complied with the University of the Philippines Visayas Research Ethics Boards through the approved issuance of the Student Research Ethics Clearance (UPVREB #C-6) on March 7, 2022.

REFERENCES

- Allan EL, Froneman PW, Hodgson AN. 2006. Effects of temperature and salinity on the standard metabolic rate (SMR) of the caridean shrimp *Palaemon peringueyi*. Journal of experimental marine biology and ecology. 337(1):103–108. <https://doi.org/10.1016/j.jembe.2006.06.006>
- Alonzo KHF, Cadiz RE, Traifalgar RFM, Corre Jr VL. 2017. Immune responses and susceptibility to *Vibrio parahaemolyticus* colonization of juvenile *Penaeus vannamei* at increased water temperature. Aquaculture, Aquarium, Conservation & Legislation. 10(5):1238–1247. <https://www.proquest.com/scholarly-journals/immune-responses-susceptibility-vibrio/docview/2056429332/se-2>
- AOAC. 2010. Official Methods of Analysis, 18th edition. Association of Official Analytical Chemists, Washington, DC.

- Bell TA, Lightner DV. 1988. A handbook of normal penaeid shrimp histology. Baton Rouge, LA: World Aquaculture Society.
- Cheng W, Wang LU, Chen JC. 2005. Effect of water temperature on the immune response of white shrimp *Litopenaeus vannamei* to *Vibrio alginolyticus*. *Aquaculture*. 250(3):592–601. <https://doi.org/10.1016/j.aquaculture.2005.04.060>
- Chien LT, Hwang DF, Jeng SS. 1999. Effect of Thermal Stress on Dietary Requirement of Vitamin C in Thornfish *Terapon jarbua*. *Fisheries science*. 65(5):731–735. <https://doi.org/10.2331/fishsci.65.731>
- Chim L, Lemaire P, Delaporte M, Le Moullac G, Galois R, Martin JLM. 2001. Could a diet enriched with n-3 highly unsaturated fatty acids be considered a promising way to enhance the immune defences and the resistance of Penaeid prawns to environmental stress? *Aquaculture Research*. 32(2):91–94. <https://doi.org/10.1046/j.1365-2109.2001.00534.x>
- Dawood MAO, Koshio S. 2018. Vitamin C supplementation to optimize growth, health and stress resistance in aquatic animals. *Reviews in Aquaculture*. 10(2):334–350. <https://doi.org/10.1111/raq.12163>
- de la Vega E, O'Leary NA, Shockey JE, Robalino J, Payne C, Browdy CL, Warr GW, Gross PS. 2008. Anti-lipopolysaccharide factor in *Litopenaeus vannamei* (LvALF): A broad spectrum antimicrobial peptide essential for shrimp immunity against bacterial and fungal infection. *Molecular Immunology*. 45(7):1916–1925. <https://doi.org/10.1016/j.molimm.2007.10.039>
- Destoumieux-Garzón D, Rosa RD, Schmitt P, Barreto C, Vidal-Dupiol J, Mitta G, Gueguen Y, Bachere E. 2016. Antimicrobial peptides in marine invertebrate health and disease. *Philosophical Transactions of the Royal Society B: Biological Sciences*. 371(1695):20150300. <https://doi.org/10.1098/rstb.2015.0300>
- Díaz F, Re AD, Sánchez A, Cruz H, González RA, Sanchez LN, Licea A, Ponce-Rivas E, Muñoz-Márquez ME, Giffard I, Rosas C. 2013. The effects of exposure to critical thermal maxima on the physiological, metabolic, and immunological responses in adult white shrimp *Litopenaeus vannamei* (Boone). *Marine and Freshwater Behaviour and Physiology*. 45(6):365–374. <https://doi.org/10.1080/10236244.2013.771911>
- Ebadi H, Zakeri M, Mousavi SM, Yavari V, Souri M. 2021. The interaction effects of dietary lipid, vitamin E and vitamin C on growth performance, feed utilization, muscle proximate composition and antioxidant enzyme activity of white leg shrimp (*Litopenaeus vannamei*). *Aquaculture Research*. 52(5):2048–2060. <https://doi.org/10.1111/are.15056>
- Fachri M, Supriana T, Baharuddin M. 2021. Analysis of affecting factors of Vannamei shrimp (*Litopenaeus Vannamei*) production farming. In: *IOP Conference Series: Earth and Environmental Science*. 782(2):022020. <https://dx.doi.org/10.1088/1755-1315/782/2/022020>
- FAO. 2020. The state of world fisheries and aquaculture 2020: Sustainability in action. Food and Agriculture Organization of the United Nations. Rome. www.fao.org/
- Fracalossi DM, Allen ME, Nichols DK, Oftedal OT. 1998. Oscars, *Astronotus ocellatus*, have a dietary requirement for vitamin C. *The Journal of nutrition*. 128(10):1745–1751. <https://doi.org/10.1093/jn/128.10.1745>
- Genio JE, Traifalgar RF, Corre Jr V. 2015. Application frequency of dietary *Vibrio harveyi* lipopolysaccharide (LPS) on growth and white spot syndrome virus resistance of post larvae *Penaeus monodon*. *The Israeli Journal of Aquaculture-Bamidgeh*. 67. <https://doi.org/10.46989/001c.20717>
- Gong H, Jiang DH, Lawrence A, Gatlin III D. 2001. Lipid nutrition of *Litopenaeus vannamei* - Responsible Seafood Advocate. Global Seafood Alliance. [accessed 2024 Jul 1]. <https://www.globalseafood.org/advocate/lipid-nutrition-of-litopenaeus-vannamei/>
- González-Félix ML, Gatlin III DM., Lawrence AL, Perez-Velazquez M. 2002. Effect of various dietary lipid levels on quantitative essential

- fatty acid requirements of juvenile Pacific white shrimp *Litopenaeus vannamei*. Journal of the World Aquaculture Society. 33(3):330–340. <https://doi.org/10.1111/j.1749-7345.2002.tb00509.x>
- Guo B, Wang F, Dong S, Dong Y, Tian X. 2010. The effects of cyclical temperature changes on growth and physiological status of *Litopenaeus vannamei*. Aquaculture international. 18:921–932. <https://doi.org/10.1007/s10499-009-9314-y>
- Harrison FE, May JM. 2009. Vitamin C function in the brain: vital role of the ascorbate transporter SVCT2. Free Radical Biology and Medicine. 46(6):719–730. <https://doi.org/10.1016/j.freeradbiomed.2008.12.018>
- Hernández-López J, Gollas-Galván T, Vargas-Albores F. 1996. Activation of the prophenoloxidase system of the brown shrimp *Penaeus californiensis* (Holmes). Comparative Biochemistry and Physiology Part C: Pharmacology, Toxicology and Endocrinology. 113(1):61–66. [https://doi.org/10.1016/0742-8413\(95\)02033-0](https://doi.org/10.1016/0742-8413(95)02033-0)
- Huang F, Jiang M, Wen H, Wu F, Liu W, Tian J, Shao H. 2016. Dietary vitamin C requirement of genetically improved farmed Tilapia, *Oreochromis niloticus*. Aquaculture Research. 47(3):689–697. <https://doi.org/10.1111/are.12527>
- Janewanthanakul S, Supungul P, Tang S, Tassanakajon A. 2020. Heat shock protein 70 from *Litopenaeus vannamei* (LvHSP70) is involved in the innate immune response against white spot syndrome virus (WSSV) infection. Developmental & Comparative Immunology. 102: 103476. <https://doi.org/10.1016/j.dci.2019.103476>
- Jia X, Wang F, Lu Y, Zhang D, Dong S. 2014. Immune responses of *Litopenaeus vannamei* to thermal stress: a comparative study of shrimp in freshwater and seawater conditions. Marine and Freshwater Behaviour and Physiology. 47(2):79–92. <https://doi.org/10.1080/10236244.2014.894349>
- Joseph A, Philip R. 2007. Acute salinity stress alters the hemolymph metabolic profile of *Penaeus monodon* and reduces immunocompetence to white spot syndrome virus infection. Aquaculture. 272:87–97.
- Junprung W, Supungul P, Tassanakajon A. 2017. HSP70 and HSP90 are involved in shrimp *Penaeus vannamei* tolerance to AHPND-causing strain of *Vibrio parahaemolyticus* after non-lethal heat shock. Fish & Shellfish Immunology. 60:237–246. <https://doi.org/10.1016/j.fsi.2016.11.049>
- Junprung W, Supungul P, Tassanakajon A. 2021. Structure, gene expression, and putative functions of crustacean heat shock proteins in innate immunity. Developmental & Comparative Immunology. 115:103875. <https://doi.org/10.1016/j.dci.2020.103875>
- Junprung W. 2019. Functional characterization of heat shock proteins during stress response in Pacific white shrimp *Penaeus vannamei* and brine shrimp *Artemia franciscana* [PhD Thesis]. Ghent University.
- Kong F, Zhu Y, Yu H, Wang X, Abouel Azm FR, Yuan J, Tan Q. 2021. Effect of dietary vitamin C on the growth performance, non-specific immunity and antioxidant ability of red swamp crayfish (*Procambarus clarkii*). Aquaculture. 541:736785. <https://doi.org/10.1016/j.aquaculture.2021.736785>
- Kumar N, Chandan NK, Gupta SK, Bhushan S, Patole PB (2022) Omega-3 fatty acids effectively modulate growth performance, immune response, and disease resistance in fish against multiple stresses. Aquaculture. 547:737–506. <https://doi.org/10.1016/j.aquaculture.2021.737506>
- Lee MH, Shiau SY. 2002. Dietary vitamin C and its derivatives affect immune responses in grass shrimp, *Penaeus monodon*. Fish & Shellfish Immunology. 12(2):119–129. <https://doi.org/10.1006/fsim.2001.0357>
- Liao IC, Chien YH. 2011. The Pacific White Shrimp, *Litopenaeus vannamei*, in Asia: The World's Most Widely Cultured Alien Crustacean. In: Galil BS, Clark PE, Carlton JT, editors. In the Wrong Place - Alien Marine Crustaceans: Distribution, Biology and Impacts. Dordrecht: Springer Netherlands. pp. 489–519. https://doi.org/10.1007/978-1-4020-9888-8_17

- org/10.1007/978-94-007-0591-3_17
- Livak KJ, Schmittgen TD. 2001. Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the $2^{-\Delta\Delta CT}$ Method. *Methods*. 25(4):402–408. <https://doi.org/10.1006/meth.2001.1262>
- Maliwat GC, Velasquez S, Robil JL, Chan M, Traifalgar RF, Tayamen M, Ragaza JA. 2017. Growth and immune response of giant freshwater prawn *Macrobrachium rosenbergii* (De Man) postlarvae fed diets containing *Chlorella vulgaris* (Beijerinck). *Aquaculture Research*. 48(4):1666–1676. <https://doi.org/10.1016/j.aaf.2020.07.00>
- Matesanz N, Park G, McAllister H, Leahey W, Devine A, McVeigh GE, Gardiner TA, McDonald DM. 2010. Docosahexaenoic acid improves the nitroso-redox balance and reduces VEGF-mediated angiogenic signaling in microvascular endothelial cells. *Investigative ophthalmology & visual science*. 51(12):6815–6825. <https://doi.org/10.1167/iovs.10-5339>
- Mercier FE, Ragu C, Scadden DT. 2012. The bone marrow at the crossroads of blood and immunity. *Nature Reviews Immunology*. 12(1):49–60. <https://doi.org/10.1038/nri3132>
- Mercier L, Racotta IS, Yepiz-Plascencia G, Muhlia-Almazán A, Civera R, Quiñones-Arreola MF, Palacios E. 2009. Effect of diets containing different levels of highly unsaturated fatty acids on physiological and immune responses in Pacific whiteleg shrimp *Litopenaeus vannamei* (Boone) exposed to handling stress. *Aquaculture Research*. 40(16):1849–1863. <https://doi.org/10.1111/j.1365-2109.2009.02291.x>
- Millward DJ. 2012. Nutrition and sarcopenia: evidence for an interaction. *Proceedings of the nutrition society*. 71(4):566–575. <https://doi.org/10.1017/S0029665112000201>
- Ming J, Xie J, Xu P, Ge X, Liu W, Ye J. 2012. Effects of emodin and vitamin C on growth performance, biochemical parameters and two HSP70s mRNA expression of Wuchang bream (*Megalobrama amblycephala* Yih) under high temperature stress. *Fish & Shellfish Immunology*. 32(5):651–661. <https://doi.org/10.1016/j.fsi.2012.01.008>
- Montoya N, Molina C. 1995. Optimun Supplemental Level of L-Ascorbyl-2-Phosphate-Mg to Diet for White Shrimp *Penaeus vannamei*. *Fisheries science*. 61(6):1045–1046. https://www.jstage.jst.go.jp/article/fishsci1994/61/6/61_6_1045/_article
- Muthaiyan R, Krishnan S, Periyasamy S, Chetri Z, Nandakumar R. 2020. Immune response of shrimp *Peneaus monodon* against *Vibrio parahaemolyticus*. *World News of Natural Sciences*. 30(2):150–168. <http://agro.icm.edu.pl/agro/element/bwmeta1.element.agro-b268f7a7-c214-4154-b001-e665fa775992>
- Nakano T, Kameda M, Shoji Y, Hayashi S, Yamaguchi T, Sato M. 2014. Effect of severe environmental thermal stress on redox state in salmon. *Redox Biology*. 2:772–776. <https://doi.org/10.1016/j.redox.2014.05.007>
- Ndong D, Chen YY, Lin YH, Vaseeharan B, Chen JC. 2007. The immune response of tilapia *Oreochromis mossambicus* and its susceptibility to *Streptococcus iniae* under stress in low and high temperatures. *Fish & Shellfish Immunology*. 22(6):686–694. <https://doi.org/10.1016/j.fsi.2006.08.015>
- Niu J, Liu Y-J, Tian L-X, Mai K-S, Lin H-Z, Chen X, Yang H-J, Liang G-Y. 2011. Influence of dietary phospholipids level on growth performance, body composition and lipid class of early postlarval *Litopenaeus vannamei*: Body composition and lipid class of early postlarval *Litopenaeus vannamei*. *Aquaculture Nutrition*. 17(2):e615–e621. <https://doi.org/10.1111/j.1365-2095.2010.00807.x>
- Niu J, Tian LX, Liu YJ, Mai KS, Yang HJ, Ye CX, Gao W. 2009. Nutrient values of dietary ascorbic acid (L-ascorbyl-2-polyphosphate) on growth, survival and stress tolerance of larval shrimp, *Litopenaeus vannamei*. *Aquaculture Nutrition*. 15(2):194–201. <https://doi.org/10.1111/j.1365-2095.2008.00583.x>
- Pan LQ, Hu FW, Jing FT, Liu HJ. 2008. The effect of different acclimation temperatures on the prophenoloxidase system and other defence

- parameters in *Litopenaeus vannamei*. *Fish & Shellfish Immunology*. 25(1):137–142. <https://doi.org/10.1016/j.fsi.2008.03.016>
- Patel KV, Mohanty JG, Kanapuru B, Hesdorffer C, Ershler WB, Rifkind JM. 2013. Association of the red cell distribution width with red blood cell deformability. In: *Oxygen Transport to Tissue XXXIV*. Springer. p. 211–216. https://doi.org/10.1007/978-1-4614-4989-8_29
- Rahman MM, Corteel M, Dantas-Lima JJ, Wille M, Alday-Sanz V, Pensaert MB, Sorgeloos P, Nauwynck HJ. 2007. Impact of daily fluctuations of optimum (27 °C) and high water temperature (33 °C) on *Penaeus vannamei* juveniles infected with white spot syndrome virus (WSSV). *Aquaculture*. 269(1):107–113. <https://doi.org/10.1016/j.aquaculture.2007.04.056>
- Roberts RJ, Agius C, Saliba C, Bossier P, Sung YY. 2010. Heat shock proteins (chaperones) in Fish and shellfish and their potential role in relation to fish health: a review. *Journal of Fish Diseases*. 33(10):789–801. <https://doi.org/10.1111/j.1365-2761.2010.01183.x>
- Sabry-Neto H, Nunes AJP. 2015. Performance and immunological resistance of *Litopenaeus vannamei* fed α -1,3/1,6-glucan-supplemented diet after per os challenge with the Infectious myonecrosis virus (IMNV). *Revista Brasileira de Zootecnia*. 44:165–173. <https://doi.org/10.1590/S1806-92902015000500001>
- Traifalgar RE, Kira H, Thanh Tung HA, Raafat ME, Laining A, Yokoyama S, Ishikawa M, Koshio S, Serrano AE, Corre V. 2010. Influence of dietary fucoidan supplementation on growth and immunological response of juvenile *Marsupenaeus japonicus*. *Journal of the World Aquaculture Society*. 41:235–244. <https://doi.org/10.1111/j.1749-7345.2010.00363.x>
- Van Gaal LE, De Leeuw IH. 1989. Reverse T₃ changes during protein supplemented diets. Relation to nutrient combustion rates. *Journal of endocrinological investigation*. 12(1989):799–803. <https://doi.org/10.1007/BF03350064>
- Vargas-Albores F. 1995. The defense system of brown shrimp (*Penaeus californiensis*): Humoral recognition and cellular responses. *Journal of marine biotechnology*. 3(1):153–156.
- Wendelaar-Bonga SE. 1997. The stress response in fish. *Physiological Reviews*. 77(3):591–625. <https://doi.org/10.1152/physrev.1997.77.3.591>
- Yan J, Liang X, Zhang Y, Li Y, Cao X, Gao J. 2017. Cloning of three heat shock protein genes (HSP70, HSP90 α and HSP90 β) and their expressions in response to thermal stress in loach (*Misgurnus anguillicaudatus*) fed with different levels of Vitamin C. *Fish & Shellfish Immunology*. 66:103–111. <https://doi.org/10.1016/j.fsi.2017.05.023>
- Zehra S, Khan MA. 2012. Dietary Vitamin C Requirement of Fingerling, *Cirrhinus mrigala* (Hamilton), Based on Growth, Feed Conversion, Protein Retention, Hematological Indices, and Liver Vitamin C Concentration. *Journal of the World Aquaculture Society*. 43:648–658. <https://doi.org/10.1111/j.1749-7345.2012.00597.x>
- Zhang X, Yuan J, Sun Y, Li S, Gao Y, Yu Y, Liu C, Wang Q, Lv X, Zhang X, Ma KY, Wang X, Lin W, Wang L, Zhu X, Zhang C, Zhang J, Jin S, Yu K. ... Xiang J. 2019. Penaeid shrimp genome provides insights into benthic adaptation and frequent molting. *Nature Communications*. 10(1):356. <https://doi.org/10.1038/s41467-018-08197-4>
- Zhou QC, Li CC, Liu CW, Chi SY, Yang QH. 2007. Effects of dietary lipid sources on growth and fatty acid composition of juvenile shrimp, *Litopenaeus vannamei*. *Aquaculture Nutrition*. 13(3):222–229. <https://doi.org/10.1111/j.1365-2095.2007.00470.x>

