

SHORT COMMUNICATION

Shell Dimension-Weight Relationship and Condition Index of the Invasive Mussel *Mytella strigata* in Sasmuan, Pampanga, Philippines

Dante M. Mendoza^{1*} , Michelle Grace B. Aquino², and Gerondina C. Mendoza¹

¹Department of Fisheries Science, College of Agriculture Systems and Technology, Pampanga State Agricultural University, Magalang, Pampanga, Philippines

²Department of Aquaculture, College of Fisheries, Central Luzon State University, Science City of Muñoz, Nueva Ecija, Philippines

ABSTRACT

A total of 300 *Mytella strigata* were collected at three sampling points in the mussel coastal demonstration farm in Sasmuan, Pampanga, and subjected to morphometric and gravimetric measurements for the evaluation of its growth pattern and condition index. Designated sampling sites were based on the distances to the river mouth (200 m, 400 m, and 600 m) with varying water quality parameters (temperature, salinity, pH, water velocity, and chlorophyll-a). Mussels in Site 2 had significantly larger shell dimensions (SL = 2.862 cm, SW = 1.458 cm, SH = 0.926 cm), shell weight (0.640 g), and total live weight (1.150 g). Significantly higher ($p < 0.05$) soft-tissue dry weight (0.083 g), however, was obtained in Site 3, while comparable ($p > 0.05$) soft tissue weight was observed in Site 2 (0.364 g) and Site 3 (0.413 g). Nevertheless, mussels in all sites demonstrated negative allometric growth ($b < 3$). The length-weight relationship revealed a weak to moderate, positive correlation. A significantly higher ($p < 0.05$) condition index was obtained in Site 3, which could have been influenced by higher chlorophyll-a and substantially lower water velocity. As such, this signifies its potential to have a well-established population in this spatial distance (600 m) from the river mouth. Hence, regular harvest and product development is recommended as strategic options to control its proliferation.

*Corresponding Author: dante_mendoza@psau.edu.ph

Received: July 4, 2022

Accepted: January 18, 2023

Keywords: *M. strigata*, shell dimension-weight relationship, condition index

Invasive alien species are considered major drivers of biodiversity loss worldwide (Convention on Biological Diversity 2016). Their establishment in a certain environment may implicate long-term direct and indirect impacts such as break down of biogeographic realms, decline in the abundance of native population leading to possible extinction, alteration of trophic networks, affecting fishery catches, and adversely impacting both shellfish and mangrove crab farming (Pysek et al. 2020; Joshi 2006; Fabiosa et al. 2021; Fuertes et al. 2021). In contrast, introducing non-native species also had positive impacts, including food items and livelihood opportunities to the local communities, aquaculture, and for culture and fisheries enhancement (Fuertes et al. 2021; Guerrero 2014).

In recent years, *Mytella strigata* (primarily referred to as *M. charruana* by Rice et al. 2016; Mediodia

et al. 2017; Vallejo et al. 2017; Parana et al. 2019) as identified by Fuertes et al. (2021) is one of the reported invasive species that was introduced in the Philippines (Vallejo et al. 2019). This species was first reported in Manila Bay in 2014 (Rice et al. 2016) and later found to be distributed widely in the neighboring provinces like Pangasinan, Bataan, Cavite, Bulacan, Cagayan, La Union, Ilocos Sur, Zambales, and Pampanga (Parana et al. 2019; Rice et al. 2016; Fuertes et al. 2021). In addition, the species' occurrence was also recorded in Panguil Bay, Southern Philippines (Fabiosa et al. 2021). The species' potential invasiveness is reflected in its tolerance to various environmental conditions (Mediodia et al. 2017).

Currently, most of the studies on *M. strigata* in the country mainly focused on its morphological description, distribution, and invasion of the Philippine waters. Assessment of its biological condition like

morphometric relationship, growth pattern, and condition index are still limited. The length-weight relationship is essential for the assessment of growth and estimation of biomass production of species (Park and Oh 2002; Aban et al. 2017). On the other hand, studying bivalve growth generates information on the environment's quality, particularly temperature and food availability patterns, as depicted in several studies (Thebault et al. 2008; Laing 2000; Mann et al. 2013). Further, the selection of site suitability for bivalve mariculture can be potentially associated with the condition index of the present animal in the area (Sasikumar and Krishnakumar 2011). The condition index is widely used as a parameter to determine the ecological and physiological status of a particular species (Zeng and Yang 2022). Moreover, the intensity of bivalve aquaculture in an area can be predicted using the condition index as an indicator (Filgueira et al. 2013). From a commercial viewpoint, the index relates the proportion of bivalve meat relative to its shell's weight, which is crucial in assessing quality and marketability (Župan and Šarić 2014). Hence, a low condition index suggests the unsuitability of the bivalve for harvest (Ezgeta-Balić et al. 2012). This parameter is commonly associated with reproductive development and food availability (Mladineo et al. 2007; Filgueira et al. 2013). The condition index increases during gonad maturation and decreases during spawning (Duisan et al. 2021).

Sasmuan is a coastal municipality of Pampanga situated along the coastline of Manila Bay. The town is known as the home of the newly conferred Sasmuan Bangkung Malapad Critical Habitat and Ecotourism Area (SBMCHEA) (DENR Region 3 2021). Recently, the local government established a demonstration farm for green mussels (*P. viridis*) in the area that will provide an adequate supply of seeds for restocking and transplantation. However, *M. strigata* was also observed to exist in the estuarine waters of Sasmuan. As this species is categorized as invasive and known to affect shellfish farming, this study was conducted to determine the important water quality variables contributing to its proliferation in the area. Specifically, the length-weight relationship and condition index were identified as biological indicators.

Mussel samples were collected in three established sampling points in the Technology Demonstration Farm of the Bureau of Fisheries and Aquatic Resources (BFAR) Regional Field Office 3 and the Local Government Unit (LGU) of Sasmuan, Pampanga (Figure 1). The designated collection area was based on the distance of culture lines from the mouth of Pasak River, namely Site 1 (200 m), Site 2 (400 m), and Site 3 (600 m).

In-situ assessments of water quality parameters such as temperature, salinity, and pH were evaluated using portable pen-type monitoring devices



Figure 1. Map of the sampling sites (extracted from Google Earth, 2022).

(RCYAGO), while water velocity was measured with the float method. A constant volume of 100 mL of water sample was collected in the area and subjected to chlorophyll-a analysis through acetone extraction (90%) and spectrophotometric approach. The total chlorophyll-a concentration was calculated using the formula from Aminot and Rey (2001):

$$\text{Chl. a} = \frac{(11.85 \times (E_{664} - E_{750}) - 1.54 \times (E_{647} - E_{750}) - 0.08 \times (E_{630} - E_{750})) \times V_e}{L \times V_f}$$

where L is the cuvette light path (cm), V_e is the extraction volume (ml), V_f is the filtered volume (l), and Chl. a is the chlorophyll a concentration (mg m^{-3}). Each measurement of water parameters was replicated five times.

Specimens of *M. strigata* (100 per site) were randomly taken by a commissioned fisher from the culture lines. Live mussel samples were transported to the laboratory, cleaned with continuous tap water, and manually removed fouling organisms. Morphometric measurements (SL = shell length, SW = shell width, and SH = shell height) and the total live weight were taken using a digital caliper with 0.1 mm sensitivity and digital weighing balance with 0.01 g sensitivity, respectively (Figure 2). Live mussels were then subjected to freezing temperatures for 24 hours and thawed in tap water for two hours. Soft tissue was




Figure 2. Illustration of dimensions in the shell of *M. strigata*.

removed using a sterile scalpel. The shell and soft tissue were blotted in tissue paper and air dried for an hour. Air-dried shell and soft tissue samples were put in a preheated oven and dried at 60°C for one hour until a constant weight was obtained. Weight measurements (SDW = shell dry weight, STW = soft tissue weight) were then measured using a similar weighing balance.

Simple linear regression was employed to evaluate the relationships between shell dimensions (SL, SW, and SH) and total weight. Condition index was determined using the formula of Davenport and Chen (1987):

$$\text{CI} = (\text{dry soft tissue weight} / \text{dry shell weight}) \times 100\%$$

Data on the water quality parameters, shell dimensions, weight measurements, and condition index were subjected to a one-way Analysis of Variance (ANOVA). In addition, Duncan's Multiple Range Test (DMRT) was used to determine the significant differences among means of the mussel's biometric data (shell length, shell width, shell height, live weight, dry shell weight, soft tissue weight, and soft tissue dry weight) and condition index from different sampling sites.

Water quality parameters across sampling sites showed significant variations (Table 1). In the present study, temperature readings also showed significant variations based on spatial differences from the river mouth. Significant differences may associate directly with the velocity of the water (Sinokrot and Gulliver 2000). The salinity level in the coastal demonstration farm also demonstrated significant differences, where site 3 had the highest salinity concentration compared to sites 1 and 2. Nevertheless, both temperature and salinity levels across sites conform to the mean readings of Huang et al. (2021), where a higher prevalence rate (> 70%) of *M. strigata* was observed. In addition, the recorded pH levels among sites were also within the pH range of 7.51 – 8.07, as described in the study of Mediodia et al. (2017). However, significant variations in the rate of water velocity may be linked to the distances of the sites and the force of the outgoing water. This may suggest that the closer to the river mouth, the faster

Table 1. Mean readings of physico-chemical parameters in the three sampling sites.

Sampling Site	Temperature (°C)	Salinity (ppt)	pH	Water Velocity (cm s-1)	Chlorophyll-a (mg m-3)
1	28.44 ^b	21.00 ^b	7.51	19.32 ^a	0.428 ^b
2	29.42 ^a	21.14 ^b	7.52	14.58 ^b	0.699 ^a
3	29.40 ^a	21.96 ^a	7.51	10.51 ^c	0.766 ^a

Values are expressed as mean ± SD. Means in the same column having different superscripts are significantly different ($p < 0.05$).

the rate of water velocity. Further significant variations in chlorophyll-a concentration between sampling points may be associated with the water's velocity rate. Pan et al. (2009) claimed that water velocity in the lotic region affects the nutrient concentration and increases the rate of suspended solids resulting in limitation on algal growth. However, the site of the present study demonstrated a lower chlorophyll-a level, and it is below the recommended minimum concentration of 1 mg m⁻³ for the potential bivalve site (Saxby 2002).

There are significant variations between sites on the morphometric and weight measurements of the mussels (Table 2). The shell dimensions (SL, SW, and SH), total weight, and shell dry weight were significantly higher in site 2. The highest STW was significantly higher in sites 3 (0.413±0.239) and 2 (0.364±0.191) compared to site 1 (0.304±0.180). The soft tissue dry weight (STDW), however, was observed to be significantly higher in site 3. Existed differences may be attributed to the surrounding

waters' environmental conditions among sampling sites, particularly with temperature, salinity, and chlorophyll-a. Temperature is considered a critical parameter with a pronounced effect on chemical processes (i.e., salinity) and has the strongest effect in driving plankton abundance (Boyd and Lichtkoppler 1979; Rasconi et al. 2017). These parameters were also observed among physico-chemical factors that influenced the soft tissue weight of the mussels *Perna perna* and *Mytilus galloprovincialis* (Sokolowski et al. 2010; Aziz et al. 2020). In addition, mussel specimens exhibiting significantly higher total weight congruently obtained significantly higher shell dimensions and dry weight. Findings suggest that the size of the shell corresponds to the increase in the mussel's total weight and may also be associated with the higher amount of fluid it contains (Gimin et al. 2004).

Analysis of the relationship between shell dimensions and weight in Table 3 showed that the mussels among sites grow in a negative allometric

Table 2. Morphometric and weight measurements of *M. strigata*.

Measurements	Site 1	Site 2	Site 3
SL (cm)	2.723±0.336 ^b	2.862±0.343 ^c	2.556±0.290 ^a
SW (cm)	1.405±0.174 ^a	1.458±0.174 ^b	1.379±0.148 ^a
SH (cm)	0.886±0.131 ^b	0.926±0.132 ^c	0.849±0.105 ^a
TW (g)	0.872±0.282 ^a	1.150±0.331 ^c	1.021±0.284 ^b
SDW (g)	0.539±0.173 ^a	0.640±0.184 ^b	0.568±0.158 ^a
STW (g)	0.304±0.180 ^a	0.364±0.191 ^b	0.413±0.239 ^b
STDW (g)	0.062±0.036 ^a	0.074±0.039 ^a	0.083±0.048 ^b

Values are expressed as mean ± SD. Means in the same row having different superscripts are significantly different ($p < 0.05$). SL = shell length; SW = shell width; SH = shell height; TW = total weight; SDW = shell dry weight; STW = soft tissue weight; STDW = soft tissue dry weight

Table 3. Relationship of shell dimension (cm) and total weight (g) of *M. strigata*.

Relationship	n	a	b±SE	r ²	F*	Growth Pattern
Site 1 (200 m)						
SL vs TW	100	- 1.010	0.691±0.050	0.664	193.315*	(-) allometric
SW vs TW	100	- 1.129	1.424±0.082	0.754	300.723*	(-) allometric
SH vs TW	100	- 0.535	1.588±0.151	0.530	110.407*	(-) allometric
Site 2 (400 m)						
SL vs TW	100	- 0.260	0.493±0.084	0.261	34.612*	(-) allometric
SW vs TW	100	- 0.459	1.105±0.156	0.337	49.916*	(-) allometric
SH vs TW	100	0.299	0.920±0.234	0.136	15.423*	(-) allometric
Site 3 (600 m)						
SL vs TW	100	- 0.673	0.663±0.073	0.457	82.472*	(-) allometric
SW vs TW	100	- 0.636	1.201±0.151	0.392	63.079*	(-) allometric
SH vs TW	100	- 0.411	1.686±0.213	0.390	62.638*	(-) allometric

n = sample size; a = intercept; b = slope; SE = standard error; r² = coefficient of determination; * all F values are highly significant ($p < 0.01$)

pattern ($b < 3$). Results corroborate with the findings of Barros et al. (2020), Keskin et al. (2020), Parana et al. (2019), and Aban et al. (2017) on allometry relationships in different species of mussels. Hence, "b" values obtained in the present study implied that the rate of increase in length of the mussel is faster than the rate of increase in its weight. The negative allometry could be correlated with the inadequacy of suitable food in the environment (Soon et al. 2016), which was reflected in the obtained chlorophyll-a concentration. In the wild, biometric traits of marine bivalves were positively associated with the abundance of phytoplankton (Noor et al. 2021). Other factors influencing mussel shell morphology and shell growth includes growth rate, density, salinity, temperature, and water flow (Seed 1968; Lesser et al. 2010; Telesca et al. 2018; Zajac et al. 2018). Further, the length-weight relationship revealed a weak to moderate, positive correlation. Lowest coefficient of determination (r^2) values for shell dimensions (SL, SW, SH) and live weight model is lower in Site 2 compared with Sites 1 and 3. The results in the present study had lower values than the findings of Parana et al. (2019) ($r^2 = 0.811 - 0.948$). Generally, about 13.6 - 75.4% of total variations in live weight can be explained by the linear model. Thus, shell dimensions are important factors that contribute to the weight of the organism, but SH has lower reliability as predictor as it has higher rate of unexplained variation in the model. Also, mussels did not gain much weight despite increasing growth in their shell dimensions. As such, it suggests that energy is invested in the growth and strengthening of shell valves for protection against adverse environmental conditions and predators (Gimin et al., 2004).

Meanwhile, the result of the F-test indicates that all the slopes (b) of the regression lines significantly differ from zero.

In terms of condition index (CI), mussels from Site 3 had significantly higher CI (14.42%) than in Sites 1 and 2, with comparable mean CI of 12.20% and 11.17% ($p < 0.05$), respectively (Figure 3). Relative to the report of Parana et al. (2019), the obtained values are lower. Arrieche et al. (2020) claimed that a CI range of 2.01 to 13.49% may indicate that mussel reproduction is inactive. The better CI of mussels in site 3 could be linked to relatively higher chlorophyll-a concentration, significantly slower water velocity, and higher salinity level in the area compared with other sites. The chlorophyll-a estimates phytoplankton biomass (Flemer 1969; Pan et al. 2009). Phytoplankton serves as a primary component of the diet of filter-feeders like mussels (Hamli et al. 2020). Meanwhile, excessive current speed primarily inhibits the mussel's feeding activity, resulting in a negative influence on the CI (Noor et al. 2021). CI and salinity may associate with the clearance rate of mussel. The clearance rate is the volume of water cleared of suspended particles per unit of time (Riisgard 2001). For instance, Vlastic et al. (2018) observed that the clearance rate of the bearded horse mussel *Modiolus barbatus* increases with increasing salinity reaching a maximum of $4.23 \text{ L h}^{-1} \text{ g}^{-1}$ at 25 psu and then decreases when salinity concentration exceeds 25 psu. Further, Celik (2012) claimed that there is a variation in the CI of mussels according to the location in which environmental parameters significantly differ. The obtained CI values in the present study may also relate to the reproductive stage of the mussel. According to Hamli et al. (2017),




Figure 3. Mean condition index of *bahong* collected in three sampling locations in the Mussel Coastal Demonstration Farm of Sasmuan, Pampanga.

the CI increases as the gonad reaches the maturation stage, and it decreases when the gonads are at the spent and resting phase. Therefore, mussels across the sites could be characterized as an adult and are already reproductively mature based on the measured shell lengths (Stenyakina et al. 2010; Lim et al. 2018; Lodeiros et al. 2021).

This study concludes that there were significant variations in the physico-chemical conditions in the three sites. The current environmental conditions in Site 3 are favorable to support the growth of *M. strigata*, as evident by the result of morphometrics and condition index. The present findings can be used as an essential reference for developing effective monitoring and management strategies to counter the invasive nature of the species in the area and eliminate its threat against the farming of green mussels. However, a study that will emphasize the extent of its invasiveness and economic impact on green mussel farming must be investigated to identify appropriate policies to secure the viability of the existing green mussel farming in the area. Moreover, it is recommended for future studies to focus on the improved utilization of the invasive species, of which the present findings can be used as a baseline. Among the potential strategies to control its population is to use the species for developing and commercializing food and non-food products. However, studies with a broader scope that will capture spatial and temporal evaluation of the overall condition of its population in the coastal waters of Pampanga and nearby areas are indispensable to determining its viability to support such undertakings.

ACKNOWLEDGMENT

The authors would like to express their gratitude to the Municipal Agriculture Office of Sasmuan, Pampanga and BFAR Regional Field Office 3 for allowing the researchers to conduct the said study. Also, special thanks to the commissioned boatman and fisher who assisted the researchers in collecting specimens.

AUTHOR CONTRIBUTION

Mendoza DM: Conceptualization, Methodology, Investigation, Formal Analysis, Writing - Review and Editing. **Aquino MGB:** Methodology, Formal Analysis, Writing - Review and Editing. **Mendoza GC:** Methodology, Formal Analysis, Writing - Review and Editing.

CONFLICTS OF INTERESTS

The authors declare that there is no conflict of interest in the publication of the research findings.

ETHICS STATEMENT

No animal was used or subjected by the authors to experimental studies.

REFERENCES

- Aban SM, Argente FAT, Raguindin, RS, Garcia AC, Ibarra CE, and De Vera RB. 2017. Length-weight relationships of the Asian Green Mussel, *Perna viridis* (Linnaeus 1758) (Bivalvia: Mytilidae) population in Bolinao Bay, Pangasinan, Northern Philippines. *PSU Journal of Natural and Allied Sciences*. 1(1):1-6.
- Aminot A, Rey F. 2001. Chlorophyll a: Determination by spectroscopic methods. *ICES Techniques in Marine Environmental Sciences*. No. 30. pp 1-17. <http://dx.doi.org/10.25607/OBP-278>
- Arrieche D, Maeda-Martinez AN, Acosta-Balbas V, Freites L, Acosta-Salmon H, Lodeiros-Seijo C. 2020. Optimum temperature for growth of an invasive green mussel *Perna viridis* population from Venezuela, determined in an open-flow system. *Aquaculture Reports*. 16:100284. <https://doi.org/10.1016/j.aqrep.2020.100284>
- Aziz G, Layachi M, Akodad M, Baghour M, Skalli A, Moumen A. 2020. Growth weight and reproductive cycle in the mussel (*Mytilus galloprovincialis*) from Cala Iris Sea of Al Hoceima (Northern Morocco). *International Journal of Biosciences*. 16(6):140-151. <https://doi.org/10.12692/ijb/16.6.140-151>
- Barros MRE, dos Santos WJP, das Chagas RA. 2020. Morphometry and shell shape stabilization indicator (IEF) of the mussel *Mytella charruana* (d'Orbigny, 1842) (Bivalvia, Mytilidae). *Biota Amazonia* 10(1): 31-34. <http://dx.doi.org/10.18561/2179-5746/biotaamazonia.v10n1p31-34>
- Boyd CE, Lichtkoppler F. 1979. Water Quality Management in Pond Fish Culture. Research

- and Development Series No. 22. Alabama: International Center for Aquaculture Agricultural Experiment Site. 30 pp.
- Convention on Biological Diversity. 2016. Invasive Alien Species. [Accessed June 6, 2022]. <https://www.cbd.int/invasive/>
- Celik MY, Karayucel S, Karayucel I, Ozturk R, Eyuboglu B. 2012. Meat Yield, Condition Index, and Biochemical Composition of Mussels (*Mytilus galloprovincialis* Lamarck, 1819) in Sinop, South of the Black Sea. *Journal of Aquatic Food Product Technology*. 21(3):198–205. <https://doi.org/10.1080/10498850.2011.589099>
- [DENR Region 3] Department of Environment and Natural Resources Region 3. 2021. DENR declares Bangkung Malapad as Critical Habitat, Ecotourism Area. <https://r3.denr.gov.ph/index.php/news-events/press-releases/1389-denr-declares-bangkung-malapad-as-critical-habitat-ecotourism-area>
- Duisan L, Salim G, Ransangan J. 2021. Sex ratio, gonadal and condition indexes of the Asiatic hard clam, *Meretrix meretrix* in Marudu Bay, Malaysia. *Biodiversitas*. 22(11):4895–4904. <https://doi.org/10.13057/biodiv/d221123>
- Ezgeta-Balić D, Najdek M, Peharda M, Blažina M. 2012. Seasonal fatty acid profile analysis to trace origin of food sources of four commercially important bivalves. *Aquaculture*. 334–337: 89–100.
- Fabiosa MP, Abao CAA, Fabiosa NP, Uba KIN. 2021. First record of the invasive mussel *Mytella strigata* (Mollusca: Bivalvia: Mytilidae) in Panguil Bay, Southern Philippines. *Journal of Environment and Aquatic Resources*. 6:35–46. <https://doi.org/10.48031/msunjea.2021.06.03>
- Filgueira R, Comeau LA, Landry T, Grant J, Guyondet T, Mallet A. 2013. Bivalve condition index as an indicator of aquaculture intensity: A meta-analysis. *Ecological Indicators*. 25:215–229. <https://doi.org/10.1016/j.ecolind.2012.10.001>
- Flemer DA. 1969. Chlorophyll analysis as a method of evaluating the standing crop phytoplankton and primary productivity. *Chesapeake Science*. 10:301–306. <https://doi.org/10.2307/1350474>
- Fuertes VNB, de los Reyes RA, Monteclaro HM. 2021. The spread of the non-indigenous mussel species *Mytella strigata* (Hanley, 1843) in the Philippines: Ensuing issues and responses of local communities. *Regional Studies in Marine Science*. 41:101576. <https://doi.org/10.1016/j.rsma.2020.101576>
- Gimin R, Mohan R, Thinh LV, Griffiths AD. 2004. The relationship of shell dimensions and shell volume to live weight and soft tissue weight in the mangrove clam, *Polymesoda erosa* (Solander, 1786) from northern Australia. *Naga*. 27(3–4):32–35. <https://hdl.handle.net/20.500.12348/2026>
- Guerrero RD. 2014. Impacts of introduced freshwater fishes in the Philippines (1905–2013): A review and recommendations. *Philippine Journal of Science*. 143(1):49–59. https://philjournalsci.dost.gov.ph/images/pdf/pjs_pdf/vol143no1/pdf/impacts_of_introduced_freshwater_fishes_in_the_Phils.pdf
- Hamli H, Hashim N, Abdulla-Al-Asif. 2020. Isolation and potential culture of phytoplankton live feed for freshwater mussels *Sinandonta woodiana* (Lea, 1934). *Asian Journal of Animal Science*. 14(4):127–136. <https://doi.org/10.3923/ajas.2020.127.136>
- Hamli, H, Idris, MH, Rajae, AH, Kamal, AHM, Hoque, MN. 2017. Condition index of *Meretrix lyrata* (Sowerby 1851) and its relationship with water parameter in Sarawak. *Sains Malaysiana*. 46(4):545–551. <http://dx.doi.org/10.17576/jsm-2017-4604-05>
- Huang YC, Li ZK, Chen WL, Chan CC, Hsu HY, Lin, YT, Huang YS, Han, YS. 2021. First record of the invasive biofouling mussel *Mytella strigata* (Hanley, 1843) (Bivalvia: Mytilidae) from clam ponds in Taiwan. *BioInvasions Records*. 10(2):304–312. <https://doi.org/10.3391/bir.2021.10.2.08>
- Joshi RC. 2006. Invasive alien species (IAS): Concerns and status in the Philippines. In: *Proceedings of the International Workshop on the*

- Development of Database (APASD) for Biological Invasion. FFTC, Taichung, Taiwan, China, pp. 1–23.
- Keskin I, Ekici A, Serdar S. 2020. Determination of the growth performance of *Mytilus galloprovincialis* in nets at Gokceada Island. *The European Zoological Journal*. 87(1): 559–570. <https://doi.org/10.1080/24750263.2020.1818856>
- Laing I. 2000. Effect of temperature and ration on growth and condition of king scallop (*Pecten maximus*) spat. *Aquaculture*. 183(3–4):325–334. [https://doi.org/10.1016/S0044-8486\(99\)00262-8](https://doi.org/10.1016/S0044-8486(99)00262-8)
- Lesser MP, Bailey MA, Merselis DG, Morrison JR. 2010. Physiological response of the blue mussel *Mytilus edulis* to differences in food and temperature in the Gulf of Maine. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*. 156(4):541–551. <https://doi.org/10.1016/j.cbpa.2010.04.012>
- Lim JY, Tay TS, Lim CS, Lee SSC, Teo SL.-M, Tan KS. 2018. *Mytella strigata* (Bivalvia: Mytilidae): an alien mussel recently introduced to Singapore and spreading rapidly. *Molluscan Research*. 38(7):1–17. <https://doi.org/10.1080/13235818.2018.1423858>
- Lodeiros C, Hernandez-Reyes D, Salazar JM, Rey Mendez M, Gonzalez-Henriquez N. 2021. First report of the mussel *Mytella strigata* (Hanley, 1843) in the Venezuelan Caribbean from an invasion in a shrimp farm. *Latin American Journal of Aquatic Research*. 49(3):531–537. <https://doi.org/10.3856/vol49-issue3-fulltext-2626>
- Mann R, Munroe DM, Powell EN, Hofmann EE, Klinck JM. 2013. Bivalve Mollusks: Barometers of Climate Change in Arctic Marine Systems. In: Mueter FJ, Dickson DMS, Huntington HP, Irvine JR, Logerwell EA, MacLean SA, Quakenbush LT, Rosa C, editors. *Responses of Arctic Marine Ecosystems to Climate Change*. Alaska Sea Grant, University of Alaska Fairbanks. <https://doi.org/10.4027/ramecc.2013.04>
- Mediodia DP, De Leon SMS, Anasco NC, Baylon CC. 2017. Shell morphology and anatomy of the Philippine Charru Mussel *Mytella charruana* (d'Orbigny, 1842). *Asian Fisheries Science* 30:185–194. <https://doi.org/10.33997/j.afs.2017.30.3.004>
- Mladineo I, Peharda M, Orhanović S, Bolotin J, Pavela-Vrančić M, Treursić B. 2007. The reproductive cycle, condition index and biochemical composition of the horse-bearded mussel *Modiolus barbatus*. *Helgoland Marine Research*. 61:183–192. <https://doi.org/10.1007/s10152-007-0065-8>
- Noor AR, Shakil A, Hoque NF, Rahman MM, Akter S, Talukder A, Ahmad-Al-Nahid S, Wahab MA, Nahiduzzaman M, Rahman MJ, Asaduzzaman M. 2021. Effect of eco-physiological factors on biometric traits of green mussel *Perna viridis* cultured in the south-east coast of the Bay of Bengal, Bangladesh. *Aquaculture Reports*. 19:100562. <https://doi.org/10.1016/j.aqrep.2020.100562>
- Pan BZ, Wang HJ, Liang XM, Wang HZ. 2009. Factors influencing chlorophyll-a concentration in the Yangtze-connected lakes. *Fresenius Environmental Bulletin*. 18(10):1894–1900. <http://benthos.ihb.ac.cn/PanWang09FreEnvironBull.PDF>
- Parana JDV, Mejia RE, Estacio MAS, Aban SM. 2019. Density, growth pattern, and condition index of western hemisphere blue mussel (*Mytella charruana*) in the riverine system of Binmaley, Pangasinan, Philippines. *Southeast Asian Journal of Science and Technology*. 4(1):50–58.
- Park KY, Oh CW. 2002. Length-weight relationship of bivalves from coastal waters of Korea. *Naga*. 25(1):21–22. <https://hdl.handle.net/20.500.12348/2284>
- Pysek P, Hulme PE, Simberloff D, Bacher S, Blackburn TM, Carlton JT, Dawson W, Essl F, Foxcroft LC, Genovesi P, et al. 2020. Scientists' warning on invasive alien species. *Biological Reviews*. 95(6):1511–1534. <https://doi.org/10.1111/brv.12627>
- Rasconi S, Winter K, Kainz MJ. 2017. Temperature increase and fluctuation induce phytoplankton

- biodiversity loss – Evidence from a multi-seasonal mesocosm experiment. *Ecology and Evolution*. 7:2936–2946. <https://doi.org/10.1002/ece3.2889>
- Rice MA, Rawson PD, Salinas AD, Rosario WR. 2016. Identification and salinity tolerance of the western hemisphere mussel *Mytella charruana* (D'Orbigny, 1842) in the Philippines. *Journal of Shellfish Research*. 35(4):865–873. <https://doi.org/10.2983/035.035.0415>
- Riisgard HU. 2001. On measurement of filtration rates in bivalves – the stony road to reliable data: review and interpretation. *Marine Ecology Progress Series*. 211:275–291. <https://www.int-res.com/articles/meps/211/m211p275.pdf>
- Sasikumar G, Krishnakumar PK. 2011. Aquaculture planning for suspended bivalve farming systems: The integration of physiological response of green mussel with environmental variability in site selection. *Ecological Indicators*. 11(2):734–740. <https://doi.org/10.1016/j.ecolind.2010.06.008>
- Saxby SA. 2002. A review of food availability: sea water characteristics and bivalve growth performance at coastal culture sites in temperate and warm temperate regions of the world. *Fisheries Research Report*. 132:1–44. http://www.fish.wa.gov.au/Documents/research_reports/fr132.pdf
- Seed R. 1968. Factors influencing shell shape in the mussel *Mytilus edulis*. *Journal of the Marine Biological Association of the United Kingdom*. 48(3):561–584. <https://doi.org/10.1017/S0025315400019159>
- Sinokrot BA, Gulliver JS. 2000. In-stream flow impact on river water temperatures. *Journal of Hydraulic Research*. 38(5):339–349. <https://doi.org/10.1080/00221680009498315>
- Sokolowski A, Bawazir AS, Sokolowska E, Wolowicz M. 2010. Seasonal variation in the reproductive activity, physiological condition and biochemical components of the brown mussel *Perna perna* from the coastal waters of Yemen (Gulf of Aden). *Aquatic Living Resources*. 23(2):177–186. <https://doi.org/10.1051/alr/2010016>
- Soon TK., Denil DJ, Ransangan J. 2016. High mortality and poor growth of green mussels, *Perna viridis*, in high chlorophyll-a environment. *Ocean Science Journal*. 51:43–57. <https://doi.org/10.1007/s12601-016-0005-0>
- Stenyakina A, Walters LJ, Hoffman EA, Calestani C. 2010. Food availability and sex reversal in *Mytella charruana*, an introduced bivalve in the southeastern United States. *Molecular Reproduction and Development*. 77(3): 222–230. <https://doi.org/10.1002/mrd.21132>
- Telesca L, Michalek K, Sanders T, Peck LS, Thyrring J, Harper EM. 2018. Blue mussel shell shape plasticity and natural environments: a quantitative approach. *Scientific Reports*. 8:2865. <https://doi.org/10.1038/s41598-018-20122-9>
- Thebault J, Thouzeau G, Chauvaud L, Cantillanez M, Avendaño M. 2008. Growth of *Argopecten purpuratus* (Mollusca: Bivalvia) on a natural bank in Northern Chile: schrenological record and environmental controls. *Aquatic Living Resources*. 21(1):45–55. <https://doi.org/10.1051/alr:2008021>
- Vallejo Jr B, Conejar-Espedido J, Manubag L, Artiaga KCC, Damatac II AM, Imperial ICVJ, Itong TAB, Fontanilla IK, Cao EP. 2017. First record of the Charru mussel *Mytella charruana* d'Orbignyi, 1846 (Bivalvia: Mytilidae) from Manila Bay, Luzon, Philippines. *BioInvasions Records*. 6(1):49–55. <https://doi.org/10.3391/bir.2017.6.1.08>
- Vallejo BM, Aloy AB, Ocampo M, Conejar-Espedido J, Manubag LM. 2019. Manila Bay Ecology and Associated Invasive Species. In: Makowski C, Finkl C, editors. *Impacts of Invasive Species on Coastal Environments*. Coastal Research Library Vol 29. Springer, Cham. https://doi.org/10.1007/978-3-319-91382-7_5
- Vlasic M, Glavic N, Bolotin J, Hrustic E, Radic ID. 2018. Influence of salinity on physiological response of the bearded horse mussel *Modiolus barbatus* and Noah's ark shell *Arca noae*. *Ekologia (Bratislava)*. 37(4):345–357. <https://doi.org/10.2478/eko-2018-0026>

- Zajac K, Zajac T, Cmiel A. 2018. What can we infer from shell dimensions of the thick-shelled river mussel *Unio crassus*? *Hydrobiologia*. 810:415–431. <https://doi.org/10.1007/s10750-017-3098-2>
- Zeng Y, Yang H. 2022. Review of molluscan bivalve condition index calculations and application in Northern Quahogs *Mercenaria mercenaria*. *Aquaculture Research*. 52(1):23–36. <https://doi.org/10.1111/are.14866>
- Župan I, Šarić T. 2014. Growth and condition index - two important factors in mussel farming. *MESO*. 16:275–278. https://www.researchgate.net/publication/292147603_Growth_and_condition_index_-_two_important_factors_in_mussel_farming



© 2023 The authors. Published by the National Fisheries Research and Development Institute. This is an open access article distributed under the [CC BY-NC 4.0](https://creativecommons.org/licenses/by-nc/4.0/) license.